


Anatomy/physiology large bowel

Dr israa.H.Bashir
University of Pretoria



Anatomical Extent


- Terminal portion of the gastrointestinal tract extending from the ileocecal valve to the anal verge.
- Comprises the colon, rectum and anal canal.
- Average adult length: approximately 150 cm.
- Serves as the interface between the small intestine and the anorectum.

Components of the Large Bowel

- Cecum (with vermiform appendix)
- Ascending colon
- Hepatic (right colic) flexure
- Transverse colon
- Splenic (left colic) flexure
- Descending colon
- Sigmoid colon
- Rectum
- Anal canal

Distinguishing Features

- Three taeniae coli (libera, omentalis and mesocolica).
- Haustra (segmental sacculations).
- Semilunar folds (plicae semilunares coli).
- Appendices epiploicae (fat-filled peritoneal appendages).
- Straight crypts of Lieberkühn with abundant goblet cells.
- Absence of villi.
- Progressive reduction in luminal diameter from cecum to sigmoid colon.



Hallmark Microscopic Features

- Normal colon has straight, parallel crypts extending from muscularis mucosae to the luminal surface; unlike small bowel, there are no villi.
- Crypt epithelium is dominated by absorptive colonocytes and abundant MUC2-rich goblet cells; enteroendocrine and tuft cells are sparse.
- Crypt architecture is uniform: regular spacing, similar crypt diameter and orderly maturation along the crypt–surface axis.
- Lamina propria contains a physiologic mononuclear infiltrate (lymphocytes, plasma cells and macrophages); density normally increases toward the surface.
- Neutrophils in crypt epithelium (cryptitis/crypt abscesses), basal plasmacytosis, crypt distortion/branching or Paneth-cell metaplasia in the left colon support pathologic chronic-active colitis.

Colonoscopy& Clinical Relevance

- Recognition of taeniae coli, haustra and semilunar folds facilitates accurate segment identification.
- Anatomical landmarks improve cecal intubation, lesion localization and therapeutic endoscopy.
- Segment-specific anatomy influences the distribution of colorectal cancer, diverticular disease, volvulus and inflammatory bowel disease.

Cecum

- Blind-ended intraperitoneal pouch in the right iliac fossa; widest segment of the colon.
- Boundaries: ileocecal valve proximally; ascending colon distally.
- Key landmarks: ileocecal valve, appendiceal orifice, triradiate fold, convergence of the three taeniae coli.
- Relations: terminal ileum (medial), anterior abdominal wall (anterior), iliacus and psoas (posterior).
- Peritoneal relationship: usually intraperitoneal with variable mobility.
- Endoscopic recognition: appendiceal orifice + ileocecal valve confirm cecal intubation.
- Clinical relevance: sessile serrated lesions, appendiceal -orifice pathology, ileocecal Crohn disease/ TB, anomalous fixation →mobile cecum →cecal volvulus.

Ascending Colon

- Extends from the cecum to the hepatic flexure (~15-20 cm).
- Three taeniae coli, haustra and appendices epiploicae present.
- Relations: right kidney posteriorly; liver superiorly; lateral abdominal wall laterally.
- Peritoneal relationship: secondarily retroperitonea.
- Endoscopic landmarks: straight lumen with widely spaced haustra.
- Clinical relevance: right-sided CRC, angiodysplasia, important site of fluid and electrolyte absorption.

Transverse Colon

- Longest and most mobile colonic segment.
- Extends from hepatic to splenic flexure.
- Suspended by the transverse mesocolon.
- Relations: stomach superiorly; pancreas posteriorly; greater omentum anteriorly.
- Peritoneal relationship: completely intraperitoneal.
- Endoscopic relevance: redundant loops are common; requires torque reduction.
- Clinical relevance: mobile transverse mesocolon and middle colic vessels are key surgical planes. close relation to stomach and pancreas matter in locally advanced disease .

Descending Colon

- Extends from splenic flexure to sigmoid colon.
- Fixed along the left lateral abdominal wall.
- Relations: left kidney posteriorly; abdominal wall laterally.
- Peritoneal relationship: secondarily retroperitoneal.
- Endoscopic recognition: relatively fixed, narrow lumen.
- Clinical relevance: ischemic colitis ; splenic flexure is a watershed region (Griffith point). left-sided colorectal cancer.

Sigmoid Colon

- S-shaped highly mobile intraperitoneal segment extending to the rectosigmoid junction (S3).
- Attached by a long sigmoid mesocolon.
- Relations: pelvic viscera, left ureter, iliac vessels.
- Endoscopic relevance: commonest site of colonoscope loop formation.
- Clinical relevance: diverticular disease, sigmoid volvulus .

Rectum

- Measures 12-15 cm from rectosigmoid junction to anorectal junction.
- No taeniae coli, haustra or appendices epiploicae.
- Contains rectal ampulla and three transverse folds (valves of Houston).
- Surrounded by mesorectum containing vessels and lymphatics; enclosed by mesorectal fascia.
- Peritoneal relationship: upper third covered anteriorly/laterally; middle third anterior only; lower third entirely extraperitoneal.
- Relations: sacrum posteriorly; bladder/prostate (male) or uterus/vagina (female) anteriorly.
- Endoscopic landmarks: rectosigmoid junction, valves of Houston, anorectal junction.

Valves of Houston (Transverse Rectal Folds)

Definition

- Usually three permanent transverse folds of the rectum composed of mucosa, submucosa and some circular muscle.
- The middle fold (Kohlrausch's fold) is the largest and most constant.

Functions


- Support the rectal reservoir /fecal column .

Clinical Importance

- Kohlrausch's fold approximates the level of the anterior peritoneal reflection, but anatomy is variable .
- Careful endoscopic inspection is required as lesions may hide behind the folds.

Valves of Houston

(Transverse Rectal Folds)



Anal Canal

- Measures approximately 3–4 cm from anorectal junction to anal verge. entirely extraperitoneal.
- Divided into columnar, transitional and squamous zones.
- Key landmarks: anal columns, anal valves, dentate line, white line of Hilton, anal verge.
- Internal anal sphincter (smooth muscle) and external anal sphincter (striated muscle); puborectalis contributes to continence.
- Dentate line: corresponds to the endoderm-ectoderm embryological junction ;a key clinical watershed for epithelium ,sensation, vascular and lymphatic drainage .
- Endoscopic relevance: retroflexion permits assessment of distal rectum and anal canal.
- Clinical relevance: hemorrhoids, fissures, fistula-in-ano, HPV-associated anal neoplasia.

Embryologic Origin and Gut Rotation

- Primitive gut tube divides into foregut, midgut and hindgut (week 4).
- Midgut derivatives: cecum, appendix, ascending colon, proximal two-thirds of transverse colon.
- Hindgut derivatives: distal one-third of transverse colon, descending colon, sigmoid colon, rectum and upper anal canal.
- Midgut undergoes physiological herniation (week 6), 270° counter-clockwise rotation around the SMA, returning during weeks 10-12.
- Arterial territories established: SMA (midgut), IMA (hindgut).

Fixation and Anorectal Development

- Cecal bud appears during week 6; appendix develops from its distal end.
- Ascending and descending colon become secondarily retroperitoneal; transverse and sigmoid colon retain mesocolon.
- Cloaca is divided by the urorectal septum into the anorectal canal and urogenital sinus.
- Dentate line marks the transition between endoderm-derived upper anal canal and ectoderm-derived lower anal canal.
- Neural crest migration forms the enteric nervous system.


Clinical Correlation

- Hirschsprung disease: failure of neural crest migration.
- Malrotation and non-rotation result from abnormal midgut rotation.
- Mobile cecum and abnormal fixation predispose to volvulus.
- Mesenteric lymphovascular anatomy determines routes of colorectal cancer spread .

Normal Colonic Histology & Functional Microanatomy

- . Architecture: flat mucosal surface NO villi, straight, parallel crypts extend to the muscularis mucosae .
- Epithelium: Simple columnar with absorptive colonocytes + abundant goblet cells MUC2 mucus and tight junctions maintain the mucosal barrier.
- Lamina propria: normally contains plasma cells, lymphocytes, macrophages and lymphoid aggregates; epithelial neutrophils indicate active inflammation.
- Crypt-surface axis: proliferative stem/progenitor cells proliferate at the crypt base → differentiate and migrate toward the surface.
- Clinical pearl: Neutrophilic cryptitis/crypt abscesses=activity ;crypt distortion/branching + basal plasmacytosis + Paneth-cell metaplasia (distal to splenic flexure) support chronic mucosal injury.


- No villi.
- Straight, parallel crypts (of Lieberkühn).
- Abundant goblet cells.
- Thick muscularis mucosae.
- Outer longitudinal muscle condensed into teniae coli.



CLINICAL PEARL

In inflammation, crypt abscesses and architectural distortion are hallmarks of chronic colitis.

Colonic Crypt and Cells



Arterial Supply

- Superior mesenteric artery (SMA): ileocolic, right colic and middle colic arteries supply the caecum, ascending colon and proximal two-thirds of the transverse colon.
- Inferior mesenteric artery (IMA): left colic, sigmoid and superior rectal arteries supply the distal transverse colon, descending colon, sigmoid colon and upper rectum.
- Middle rectal artery (internal iliac): variable contribution to the mid rectum.
- Inferior rectal artery (internal pudendal): supplies the anal canal below the dentate line.
- Marginal artery of Drummond provides the principal longitudinal collateral arcade.
- Arc of Riolan may enlarge in chronic mesenteric occlusive disease.

Clinical Pearls

- *Griffith's point (splenic flexure) and Sudeck's point (rectosigmoid junction) are watershed areas susceptible to ischaemia.*

ARTERIAL SUPPLY


COLON, RECTUM & ANAL CANAL

From Superior Mesenteric Artery (SMA)

- Ileocolic a.
- Right colic a.
- Middle colic a.

From Inferior Mesenteric Artery (IMA)

- Left colic a.
- Sigmoid aa.
- Superior rectal a.



Venous Drainage

- Veins accompany corresponding arteries.
- SMV drains right colon and proximal transverse colon.
- IMV drains left colon and sigmoid colon, usually into the splenic vein.
- Superior rectal vein drains to the portal venous system.
- Middle and inferior rectal veins drain into the internal iliac system (systemic circulation).

Clinical Pearls

- Rectum is a major porto-systemic anastomotic site.
- Explains preferential hepatic metastases from colon cancer and dual metastatic pathways from distal rectal cancer.

Lymphatic Drainage

- Colon: epicolic → paracolic → intermediate → central (apical) lymph nodes along SMA/IMA.
- Upper rectum drains to inferior mesenteric nodes.
- Mid and lower rectum drain predominantly to internal iliac nodes.
- Anal canal above dentate line drains to internal iliac/inferior mesenteric pathways.
- Anal canal below dentate line drains to superficial inguinal lymph nodes.

Clinical Pearls


- Lymphatic spread follows arterial pedicles and determines oncological resection planes.
- Inguinal node involvement is characteristic of squamous cell carcinoma below the dentate line.

VENOUS & LYMPHATIC DRAINAGE

COLON, RECTUM & ANAL CANAL


VENOUS DRAINAGE

- SMV → Portal vein
(Right & Transverse colon)
- IMV → Splenic vein →
Portal vein
(Left colon & Sigmoid colon)
- Superior rectal vein →
IMV
- Middle rectal vein →
Internal iliac vein
- Inferior rectal vein →
Internal pudendal vein
(Systemic)



LYMPHATIC DRAINAGE

- Epicolic nodes
- Paracolic nodes
- Intermediate nodes
- Central nodes




Integrated Neural Control of the Colon

- ENS = intrinsic controller: enteric sensory neurons, interneurons and motor/secretomotor neurons generate local reflexes within the bowel wall and can function independently of the CNS.
 - Myenteric (Auerbach) plexus primarily coordinates smooth-muscle activity and motility; submucosal plexuses regulate secretion, absorption and mucosal blood flow.
 - Extrinsic autonomic input tunes ENS circuits: parasympathetic input is generally pro-mobility and pro-excitatory, whereas sympathetic pathways generally inhibit motility/secretion and regulate vascular tone.
- . Take-home: colonic function emerges from ENS-ICC-smooth-muscle integration, continuously modulated by autonomic/CNS input—not from a single neural pathway.

Enteric Nervous System

- Excitatory transmitters: acetylcholine, substance P, tachykinins.
- Inhibitory transmitters: nitric oxide, VIP, ATP/purines.

Enteric nervous system: motor and sensory organization



Myenteric (Auerbach) plexus

between circular and longitudinal muscle; principal motility plexus.

Submucosal plexuses

Meissner and Schabadasch networks regulate mucosal sensation, secretion absorption, and blood flow.

Intrinsic sensory neurons

Detect mechanical and chemical signals from the mucosa/gut wall.

Interneuronal circuits

link oral and aboral pathways to coordinate local reflexes.

Motor output

targets smooth muscle, epithelium and submucosal vasculature.


Interstitial Cells of Cajal (ICC)

- ICC are specialized mesenchymal pacemaker cells expressing KIT (CD117) and ANO1 (TMEM16A/DOG1).
- Major colonic subtypes: ICC-MY(myenteric), ICC-IM (intramuscular), ICC-SM/SMP (submucosal).
- SIP syncytium :Smooth muscle cells+ICC =PDGFR α +cells are electrically coupled and function as an integrated unit, controlling smooth muscle excitability and contractility .
- Clinical pearl: ICC network loss/ dysfunction is associated with enteric neuromuscular dysmotility , particularly slow-transit constipation and chronic intestinal pseudo-obstruction .

Extrinsic Autonomic Regulation

- Parasympathetic : Vagus nerve → proximal colon; pelvic splanchnic nerves (S2-S4) supply distal colon and rectum → ↑ motility and secretion.
- Sympathetic : thoracolumbar pathways (T10-L12) → mesenteric and hypogastric plexuses → ↓ motility and secretion, vasoconstriction, and support IAS tone.

Extrinsic Innervation of the Human Colon



Patterns of Colonic Motility & High-Resolution Colonic Manometry (HRCM)

Overview

Major motor patterns: Haustral contractions, Haustral fragmentation, Cyclic Motor Pattern (CMP), High-Amplitude Propagated Contractions (HAPCs), Simultaneous Pressure Waves (SPWs).

Haustral Contractions

Segmenting contractions, slow ,rhythmic non-propagating, that mix luminal contents, enhance water/electrolyte absorption and bacterial fermentation.(the commonest motor activity of the colon.

Haustral Fragmentation

Independent contraction of portions of a haustrum fragments stool, increasing surface area for fermentation and SCFA production.

Cyclic Motor Pattern (Rectosigmoid Brake)

Repetitive, retrograde contractions (2-6 cycles/min) in the rectosigmoid region limit rectal filling, contributing to continence.

Colonic Motor Patterns: Propulsion & Clinical Manometry

High-Amplitude Propagated Contractions (HAPCs)

Most powerful propulsive contractions (>75–100 mmHg, often >30 cm propagation). Occur after waking, after meals, and with stimulant laxatives; responsible for mass movements and defecation. Reduced in slow-transit constipation/colonic inertia.

Simultaneous Pressure Waves

Non-propulsive pressure events that are often associated with rapid movement of gas and/or liquid contents


High-Resolution Colonic Manometry (HRCM)

Uses closely spaced pressure sensors to characterize HAPCs, CMP, retrograde propagating sequences, and segmental motor patterns. Used in refractory constipation, suspected colonic inertia, pseudo-obstruction, pre-colectomy evaluation, and specialized motility assessment.

Clinical pearls


- HAPCs are the principal propulsive motor events of the colon.
- CMP functions as the rectosigmoid brake.
- HRCM is reserved for complex colonic dysmotility and selected pre-operative assessment.

High-Resolution Colonic Manometry (HRCM): Catheter Position



Illustrative catheter course through the colon

High-Amplitude Propagating Contractions (HAPCs): Manometric Pattern




Clinical pearl: HAPCs propagate aborally over long colonic segments and are the manometric correlate of colonic mass movement.

Cyclic Motor Pattern (CMP): The Rectosigmoid Brake



What the tracing shows

- Repetitive cyclic pressure activity,
- Most prominent in the rectosigmoid
- Retrograde activity limits rectal fill



High-resolution pressure topography

High resolution manometry (1cm recording site spacing)



THE DEFECATORY PROCESS

FOUR DISTINCT PHASES OF DEFECATION

1

BASAL PHASE

The rectum is largely empty; resting anal pressure exceeds rectal pressure. IAS tonic contraction (most resting anal pressure)/pelvic-floor tone, puborectalis muscle, and the rectosigmoid brake maintain continence.

2

PRE-EXPULSIVE PHASE

Propulsive colonic activity moves contents distally while the rectosigmoid brake is dampened. Rectal filling/distension evokes sensation and RAIR, allowing sampling and generating the conscious urge to defecate.

3

EXPULSIVE PHASE

The rectoanal pressure gradient reverses: IAS/pelvic floor relax reflexly and EAS/puborectalis relax voluntarily. The anal canal opens and abdominal plus rectal propulsive pressure expels stool.


4

END PHASE

After evacuation, the closing reflex restores anal sphincter and pelvic-floor tone. The anorectal angle and basal rectoanal pressure gradient are re-established, returning the system to continence.

CLINICAL PEARL Dyssynergic defecation reflects failure of coordinated rectal propulsion with puborectalis/EAS relaxation during the expulsive phase; anorectal manometry plus balloon-expulsion testing assesses this physiology.

Defecography: Mechanics of Defecation



During evacuation, the anorectal angle widens, the anal canal opens, and the pelvic floor/perineum descends, permitting expulsion of rectal contrast.

Reflexes

- Gastrocolic, colocolonic and rectocolic reflexes coordinate transit.
- Rectoanal inhibitory reflex (RAIR): Rectal distension→ transient IAS relaxation allowing anorectal sampling, continence is preserved by the EAS/ puborectalis until defecation is appropriate. it is mediated predominantly by enteric inhibitory neurons (NO/VIP).
- Defecation reflex: Rectal distension→ IAS relaxation+rectal propulsion; coordinated voluntary relaxation of the puborectalis and EAS permits evacuation

Key Clinical Pearls

- The colonic motility is an integrated neuromuscular process , Not simply smooth- muscle contraction .
- The SIP syncytium (SMC-ICC-PDGFR α + cells) Integrates neural signals into coordinated motor activity.
- HRCM identifies distinct motor patterns particularly HAPCs and cyclic motor patterns .

Clinical Correlations

- Hirschsprung disease: congenital distal aganglionosis and absent RAIR.
- Slow-transit constipation: ↓HAPCs ± ICC/ enteric neuromuscular dysfunction.
- Defecatory disorders: impaired rectoanal coordination→ obstructed evacuation.

Functional Overview

- Water and electrolyte absorption.
- Fermentation of dietary fibre by the colonic microbiota.
- Production and absorption of short-chain fatty acids.
- Formation, storage and controlled evacuation of stool.
- Maintenance of mucosal barrier and immune homeostasis.


What does the colon do?

- Fluid salvage: receives $\cong 1\text{--}2$ L/day under usual conditions ; final stool water is usually only $\sim 100\text{--}200$ mL/day.
- Electrolyte and acid-base handling: absorbs Na^+ and Cl^- , secretes HCO_3^- and K^+ , and participates in systemic potassium balance.
- Microbial fermentation: converts otherwise indigestible carbohydrate into SCFAs, gases and microbial biomass.
- Barrier and immune organ: maintains a mucus-separated microbial ecosystem while permitting sampling of antigens and mucosal tolerance.
- Storage and propulsion: dehydrates, stores and periodically expels fecal contents through coordinated segmental and mass movements.

Regional specialization along the colon


Region	Dominant physiological role
Cecum / ascending	High substrate availability; vigorous fermentation; major NaCl, SCFA and water absorption; lower pH.
Transverse	Continued fermentation and mixing ;ongoing fluid / electrolytes salvage .
Descending / sigmoid	Stool dehydration + storage; aldosterone-sensitive ENaC Na ⁺ absorption .
Rectum	Reservoir +sensory organ→ continence and coordinated defecation.

Functional compartmentalization of the colonic crypt



The colonic microbiota as an ecosystem

Colonic microbiota: spatial ecosystem and metabolic cross-feeding



Microbiota composition: what is clinically defensible?

- Scale: the colon contains the highest microbial density in the gastrointestinal tract and includes bacteria, archaea, fungi, viruses and bacteriophages.
- Dominant bacterial groups: Bacillota/Firmicutes and Bacteroidota commonly predominate, with Actinomycetota, Verrucomicrobiota and Proteobacteria/Pseudomonadota represented.
- Inter-individual variation: diet, medication, geography, age, bowel habit and sampling method strongly influence profiles.
- There is no universally “normal” microbiome, and routine commercial dysbiosis indices are not yet substitutes for validated clinical tests.

Drivers of microbiota structure and function


Driver	Physiological effect
Dietary substrate	Fiber promotes SCFA production;; protein/fat generate different microbial metabolites.
Transit time	Determine fermentation time ; slow transit→ prolonged fermentation/proteolysis, rapid transit limits microbial processing .
Oxygen / redox	Epithelial oxygen consumption favors obligate anaerobes ; inflammation favors facultative organisms.
Medications	Antibiotics, PPIs, metformin, laxatives and immunosuppressive therapy can reshape communities.
Bile acids	Primary bile acids undergo microbial deconjugation and conversion to secondary bile acids→ modulate microbiota, colonic motility, and host FXR/TGR5 signaling .
Host factors	Age, genetics, immunity, mucus and disease shape the microbial ecosystem .

Microbial metabolic functions of the colon

- Dietary fiber→ SCFAs: fermentation produces acetate, propionate and butyrate; butyrate is a major fuel for mature colonocytes and supports barrier integrity.
- SCFAs are signaling molecules: they modulate epithelial transport, mucosal immunity, enteroendocrine signaling and colonic motility.
- Microbial cross-feeding sustains the ecosystem: metabolites such as acetate and lactate can be converted by other organisms into butyrate.
- Microbes also transform bile acids, tryptophan and synthesize vitamins (K2 and several B vitamins) , generating bioactive metabolites that influence epithelial and immune function.

SCFA production and cross-feeding

Dietary fibre → microbial fermentation → host signaling



Faecal SCFA concentration is not equivalent to production: most SCFAs are rapidly absorbed and utilized.

Microbiota-immune crosstalk

- Barrier & separation – MUC2 mucus, tight junctions and antimicrobial programs restrict direct microbial contact with the epithelium.
- Controlled sensing – epithelial cells, macrophages and dendritic cells use PRRs (TLRs/NOD2) to sense microbes without excessive inflammation.
- Tolerance – secretory IgA contains luminal microbes; tolerogenic dendritic-cell/Treg pathways maintain mucosal immune homeostasis.
- Microbial metabolites – SCFAs (especially butyrate), bile-acid and tryptophan metabolites reinforce epithelial and regulatory immune pathways.
- Take-home – barrier failure + dysbiosis + maladaptive immunity can sustain IBD inflammation; routine microbiome profiling is not yet a treat-to-target endpoint.

Clinical Take-Home Framework

Translate colonic anatomy and physiology into better bedside decisions.

1 STRUCTURE → DISEASE

Regional anatomy, wall layers and vascular watershed zones explain patterns of ischaemia, inflammation, obstruction and procedural risk.

2 CRYPT → BARRIER

Stem-cell renewal, MUC2 mucus, tight junctions and mucosal immunity maintain homeostasis; persistent disruption links inflammation to dysplasia.

3 TRANSPORT → DIARRHOEA

NaCl and water absorption plus K^+/HCO_3^- handling explain the fluid, electrolyte and acid-base consequences of colonic disease.

4 MOTILITY → SYMPTOMS

ENS-ICC-smooth-muscle integration, transit and anorectal coordination provide the physiology behind constipation and defecatory disorders.

5 MICROBIOME → FUNCTION

Think metabolites rather than taxonomy: fibre-derived SCFAs—especially butyrate—support colonocyte energy, barrier integrity and immune regulation.

Last but not least: localize the physiology behind the symptom, then choose the targeted test—not the other way around.

References (1/3)

1. Standring S, ed. Gray's Anatomy: The Anatomical Basis of Clinical Practice. 43rd ed. Elsevier; 2026.
2. Chung RT, Rubin DT, Wilcox CM, eds. Sleisenger and Fordtran's Gastrointestinal and Liver Disease. 12th ed. Elsevier; 2026.
3. Sadler TW. Langman's Medical Embryology. 15th ed. Wolters Kluwer; 2023.
4. Feakins R, Borralho Nunes P, Driessen A, et al. Definitions of histological abnormalities in inflammatory bowel disease: an ECCO position paper. J Crohns Colitis. 2024;18(2):175-191. doi:10.1093/ecco-jcc/jjad142.
5. Magro F, Langner C, Driessen A, et al. European consensus on the histopathology of inflammatory bowel disease. J Crohns Colitis. 2013;7(10):827-851. doi:10.1016/j.crohns.2013.06.001.
6. Barker N, van Es JH, Kuipers J, et al. Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature. 2007;449:1003-1007. doi:10.1038/nature06196.
7. Furness JB. The enteric nervous system and neurogastroenterology. Nat Rev Gastroenterol Hepatol. 2012;9:286-294. doi:10.1038/nrgastro.2012.32.

References (2/3)

8. Sanders KM, Ward SM, Koh SD. Interstitial cells: regulators of smooth muscle function. *Physiol Rev.* 2014;94(3):859-907. doi:10.1152/physrev.00037.2013.
9. Lin AY, Du P, Dinning PG, et al. High-resolution anatomic correlation of cyclic motor patterns in the human colon: evidence of a rectosigmoid brake. *Am J Physiol Gastrointest Liver Physiol.* 2017;312(5):G508-G515. doi:10.1152/ajpgi.00021.2017.
10. Camilleri M, Ford AC, Mawe GM, et al. Chronic constipation. *Nat Rev Dis Primers.* 2017;3:17095. doi:10.1038/nrdp.2017.95.
11. Donowitz M, Ming Tse C, Fuster D. SLC9/NHE gene family, a plasma membrane and organellar family of Na⁺/H⁺ exchangers. *Mol Aspects Med.* 2013;34(2-3):236-251. doi:10.1016/j.mam.2012.05.001.
12. Alper SL, Sharma AK. The SLC26 gene family of anion transporters and channels. *Mol Aspects Med.* 2013;34(2-3):494-515. doi:10.1016/j.mam.2012.07.009.
13. Johansson MEV, Larsson JMH, Hansson GC. The two mucus layers of colon are organized by the MUC2 mucin, whereas the outer layer is a legislator of host-microbial interactions. *Proc Natl Acad Sci U S A.* 2011;108 Suppl 1:4659-4665. doi:10.1073/pnas.1006451107.
14. Parikh K, Antanaviciute A, Fawcner-Corbett D, et al. Colonic epithelial cell diversity in health and inflammatory bowel disease. *Nature.* 2019;567:49-55. doi:10.1038/s41586-019-0992-y.

References (3/3)

15. Mukhopadhye I, Louis P. Gut microbiota-derived short-chain fatty acids and their role in human health and disease. *Nat Rev Microbiol.* 2025;23:635-651. doi:10.1038/s41579-025-01183-w.
16. Fan Y, Pedersen O. Gut microbiota in human metabolic health and disease. *Nat Rev Microbiol.* 2021;19:55-71. doi:10.1038/s41579-020-0433-9.
17. Turner JR. Intestinal mucosal barrier function in health and disease. *Nat Rev Immunol.* 2009;9:799-809. doi:10.1038/nri2653.
18. Binder HJ. Role of colonic short-chain fatty acid transport in diarrhea. *Annu Rev Physiol.* 2010;72:297-313. doi:10.1146/annurev-physiol-021909-135817.
19. Sorensen MV, Matos JE, Praetorius HA, Leipziger J. Colonic potassium handling. *Pflugers Arch.* 2010;459:645-656. doi:10.1007/s00424-009-0781-9.
20. Boron WF, Boulpaep EL. *Medical Physiology.* 4th ed. Elsevier; 2022.